

Abstract Submitted to the  
International Conference on Strongly Correlated Electron Systems  
University of Michigan, Ann Arbor  
August 6-10, 2001

## **Magnetic Properties of CeNi<sub>2</sub>Ge<sub>2</sub>**

C.D.Frost<sup>1</sup>, B.D.Rainford<sup>2</sup>, F.V.Carter<sup>3</sup>, S.S.Saxena<sup>3</sup>

<sup>1</sup> *ISIS Facility, Rutherford Appleton Laboratory, OX110QX, UK*

<sup>2</sup> *Dept. of Physics, University of Southampton, SO171BJ, UK*

<sup>3</sup> *Cavendish Laboratory, Cambridge University, CB30HE, UK*

The magnetic susceptibility of the non-Fermi liquid heavy fermion compound CeNi<sub>2</sub>Ge<sub>2</sub> is highly anisotropic. The c-axis susceptibility has a broad maximum near 30K, which has previously been ascribed to short range AF correlations; at low temperatures there is a weak upturn in  $\chi_c(T)$ , while the easy axis switches from [001] to [100]. We have carried out a polarised neutron study of the induced moment in a field of 4.6T which shows that the upturn in  $\chi_c(T)$  is intrinsic. Inelastic neutron scattering revealed two components to the magnetic response, a broad crystal field excitation, centred on 25 meV, and a quasi-elastic component. Using these data we present an analysis of the single crystal susceptibility, based on a crystal field model, that describes all the main features, including the maximum in  $\chi_c(T)$  and the change in sign of the anisotropy with temperature. This model may give further insight into the origins of non-Fermi liquid behaviour in this compound.